CONTENTS

Page

Diagnosis of Tuberculosis by Complement Fixation, with Special Reference to Bovine Tuberculosis. A. EICHHORN and A. BLUMBERG 1

Availability of the Potash in Certain Orthoclase-Bearing Soils as Affected by Lime or Gypsum. LYMANN J. BRIGGS and J. F. BREAZEALE 21

Ewes' Milk: Its Fat Content and Relation to the Growth of Lambs. E. G. RITZMAN 29

Immunity Studies an Anthrax Serum. A. EICHHORN, W. N. BERG, and R. A. KELSER 37

Calcium Compounds in Soils. EDMUND C. SHOREY, WILLIAM H. FRY, and WILLIAM HAZEN 57

Studies upon the Blackleg Disease of the Potato, with Special Reference to the Relationship of the Causal Organisms. W. J. MORSE 79

Effect of Fertilizers on the Composition and Quality of Oranges. H. D. YOUNG 127

Temperature Relations of Apple-Rot Fungi. CHARLES BROOKS and J. S. COOLEY 139

Mechanism of Tumor Growth in Crowngall. ERWIN F. SMITH ... 165

Sorosporella uvella and Its Occurrence in Cutworms in America. A. T. SPEARE 189

Measurement of the Inactive, or Unfree, Moisture in the Soil by Means of the Dilatometer Method. GEORGE BOUYOUCOS 195

Leafspot-Rot of Pond Lilies Caused by Helicosporium nymphaearum. FREDERICK V. RAND 219

Studies of the Genus Phytophthora. J. ROSENBAUM 233

Tetrastichus bruchophagi, a Recently Described Parasite of Bruchophagus funebris. THEODORE D. URBAHNS 277

Arsenical Injury Through the Bark of Fruit Trees. DEANE B. SWINGLE and H. E. MORRIS 283

A Squash Disease Caused by Choanephora cucurbitarum. FREDERICK A. WOLF 319

Discovery of Internal Telia Produced by a Species of Cronartium. REGINALD H. COLLEY 329

Effect of Irrigation Water and Manure on the Nitrates and Total Soluble Salts of the Soil. F. S. HARRIS and N. I. BUTT 333

Page

The Relative Influence of Microorganisms and Plant Enzyms on the Fermentation of Corn Silage. A. R. LAMB..... 361

Little-Leaf of the Vine. FREDERIC T. BIOLETTI and LEON BONNET..... 381

Spore-Forming Bacteria of the Apiary. ARTHUR H. MCCRAY.... 399

Fusarium-Blight, or Wilt Disease, of the Soybean. RICHARD O. CROMWELL..... 421

Peanut-Wilt Caused by Sclerotium Rolfsii. J. A. MCCLINTOCK... 441

Transformation of Pseudoglobulin into Euglobulin. W. N. BERG. 449

Dissemination of the Angular Leafspot of Cotton. R. C. FAULWETTER..... 457

Index..... 477

ERRATA

Page 100, line 9, "less virulence" should read "loss of virulence."

Page 137, Table X, Plot L, "Nitrate of soda" should read "Nitrate of soda, dried blood, and sulphate of potash."

Page 164, Pl. 3, A, "250" should read "25°."

Page 198, lines 3 and 5, "B" should read "C" and "C" should read "B."

Page 198, line 9, "0.1 c. c." should read "0.01 c. c."

Pages 206, 208, Tables V and VI, "test" should read "soil."

Page 237, line 3 from bottom, "microscopic" should read "macroscopic."

Page 265, line 5 from bottom, "Pearl (22) says" should read "Pearl (21a) says".

Page 275, insert "(21a) Pearl, Raymond. Modes of Research in Genetics. 182 p., diagr. New York, 1915."

Page 275, (22), "Bul. 100" should read "Bul. 110."

Page 329, Table I, under "Host," "*Peltandra virginicia*" should read "*Peltandra virginica*."

Page 364, line 23, "(21)" should read "(5)."

Page 374, Table VIII, footnote "a" should refer to "Experiment 3" only.

Page 393, line 16, "Petite" should read "Petit."

Page 409, line 3 from bottom, "as finely granular structure as in *B. mesentericus*" should read "as finely granular in structure as *B. mesentericus*."

Page 413, line 7 from bottom, "To this, bee-larvæ bouillon," omit comma after "this."

Page 422, line 25, "(13, 14), and Manns." should read "(14), and Manns (13)."

Page 422, line 30 "(fig. 1, A-G)" should read "(fig. 1, H),"

ILLUSTRATIONS

PLATES

Temperature Relations of Apple-Rot Fungi

Page

Plate 1. A.—Grimes Golden apple affected with blackrot, showing character of the natural infections with *Sphaeropsis malorum* that were used in the temperature experiments. B.—Ben Davis apple affected with bitter-rot, showing the character of the natural infections with *Glomerella cingulata* that were used in the temperature experiments..... 164

Plate 2. Yellow Newtown apples, showing the behavior of *Sclerotinia cinerea* at different temperatures: A.—Results obtained with a culture of *S. cinerea* from a peach after two weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. B.—Results obtained with a culture of *S. cinerea* from an apple after one week at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. C.—Results obtained with a culture of *Sphaeropsis malorum* after two weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. D.—Results obtained with a culture of *Penicillium expansum* after two weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C..... 164

Plate 3. Yellow Newtown apples, showing the behavior of apple-rot organisms at different temperatures: A.—Results obtained with a culture of *Volutella fructi* after two weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. B.—Results obtained with a culture of *Glomerella cingulata* after two weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. C.—Results obtained with a culture of *Neofabraea malicorticis* after three weeks at 0°, 5°, 10°, 15°, 20°, 25°, 30° C. D.—Results obtained with a culture of *Neofabraea malicorticis* after five weeks at 0°, 5°, 10°, 15° C..... 164

Mechanism of Tumor Growth in Crowngall

Plate 4. Pith cavity of a stem of *Ricinus communis* showing proliferations of deep origin due to the injection of a mixture of strong ammonia and water.. 188

Plate 5. Opposite side of the pith cavity of *Ricinus communis* shown in Plate 4.. 188

Plate 6. 1, 2.—Sections of a green tomato fruit showing carpellary surface proliferations due to ammonium carbonate. 3.—Inner face of carpel of a green tomato fruit showing numerous mulberry-like white surface proliferations due to the injection of ammonium acetate..... 188

Plate 7. 1, 2.—Inner face of carpel wall of green tomato fruit showing proliferations due to ammonium tartrate..... 188

Plate 8. 1.—From the same series of sections as the figures on Plate 7, but showing a more copious proliferation of cells. 2.—Normal tissue of inner face of tomato carpel in vicinity of figure 1. 3.—Lining membrane of figure 2, passing into such a tumor as figure 1..... 188

Plate 9. 1, 2.—Same slide as Plate 7, but showing greater variations in the size of the proliferated cells..... 188

Plate 10. Longitudinal section of a stem of *Ricinus communis* injected with distilled water containing 10 per cent malic acid..... 188

Plate 11. 1.—Middle of an internode of *Ricinus communis* injected with 10 per cent malic acid, showing copious surface pith proliferations. 2.—Middle of a Ricinus internode injected July 18 with 10 per cent uric acid. 3.—Middle of a Ricinus internode, showing copious tumor-like proliferations due to the injection of a 20 per cent solution of urea. 4.—Two Ricinus internodes injected with a 10 per cent solution of tannic acid.... 188

Page

PLATE 12. 1, 2.—Internodes of *Ricinus communis* injected with a 5 per cent solution of sodium bicarbonate 188

PLATE 13. Internode of *Ricinus communis* showing a pith cavity occupied by numerous strands and small nodules due to injection on August 12, 1916, of a 5 per cent solution of ammonium bicarbonate 188

PLATE 14. 1.—Longitudinal section of a stem of *Ricinus communis*, showing copious proliferations due to a 5 per cent grape-sugar solution. 2.—Cross section below figure 1. 3.—Cross section of a Ricinus internode which received 0.2 c. c. of clear lime water when young 188

PLATE 15. 1, 2.—Internodes of *Ricinus communis*, showing the result of injecting a 5 per cent solution of cane sugar into young internodes of the main axis. 3.—Cross section of upper part of a young Ricinus stem showing pith cavity lined by a thin sheet of cell proliferations due to the injection of ammonium tartrate. 4.—Side view of figure 3, further enlarged 188

PLATE 16. 1.—Eight small tumors and various strands in an unopened internode of *Ricinus communis*. 2.—Section of one of the tumors shown in figure 1.. 188

PLATE 17. 1, 2.—Cross sections of two internodes of *Ricinus communis* (two plants), showing pith proliferations due to the vapors from monobasic ammonium phosphate 188

PLATE 18. 1.—Top of lower injected internode of plant 1 of *Ricinus communis*, which received 1 to 100 ammonia water. 2.—Section 1 inch below figure 1, but above the needle entrance 188

PLATE 19. 1.—Floor of upper injected internode of the main axis of plant 1 of *Ricinus communis*, showing copious proliferations from the pith. 2.—Same as figure 1, but with the stem pared down one-fourth inch 188

PLATE 20. 1.—Plant 1 of *Ricinus communis*, which received 1 to 100 ammonia water. 2.—Side view of the same 188

PLATE 21. 1.—Roof of second unopened internode below lowest injected one on main axis of plant 1 of *Ricinus communis*, which was injected with 1 to 100 ammonia water. 2.—Free-hand unstained section of tumor shown in figure 1 188

PLATE 22. 1, 2.—Plant 5 of *Ricinus communis*, injected with 1 to 1,200 ammonia water 188

PLATE 23. Plant 5 of *Ricinus communis*, branch A, showing effect of 1 to 1,200 ammonia water 188

PLATE 24. 1.—Plant 5 of *Ricinus communis*, branch B, injected with 1 to 1,200 ammonia water. 2.—Section of small marginal proliferation on wall of pith cavity shown in figure 1. 3.—Same as figure 1, but from branch C 188

PLATE 25.—Upper half of first internode above the upper injected one in the main axis of plant 5 of *Ricinus communis*, which was injected with 1 to 1,200 ammonia water 188

PLATE 26. Lower half of same internode as Plate 25, showing continuation of tumors and strands 188

PLATE 27. 1.—Cross section of a normal internode of the same age as figure 2. 2.—Main axis of plant 5 of *Ricinus communis*, injected with 1 to 1,200 ammonia water. 3.—Main axis of Ricinus plant 5, injected with 1 to 1,200 ammonia water 188

PLATE 28. 1, 2, 3, 4.—Plant 4 of *Ricinus communis*, injected with 1 to 1,200 ammonia water. Figure 3 is a ceiling 188

PLATE 29. 1–5.—Petiole and stem sections of plant 4 of *Ricinus communis*, which was injected with 1 to 1,200 ammonia water. 1.—Middle of petiole. 2, 3, 4, 5.—Parts of four unopened internodes at the top of the plant. 6.—Normal pith from Plate 24, figure 1 188

Page

PLATE 30. Ceiling of upper treated internode of plant 2 of *Ricinus communis*, which was injected with 1 to 1,200 ammonia water 188

PLATE 31. Ceiling of second (middle) injected internode of plant 3 of *Ricinus communis*, which received 1 to 1,200 ammonia water 188

PLATE 32. 1, 2.—Stem of *Ricinus communis* exposed to the vapors of monobasic ammonium phosphate (20 per cent in water) 188

PLATE 33. Stem of *Ricinus communis* in cross section showing pith-lining of normal node 188

PLATE 34. Longitudinal section of stem of *Ricinus communis* showing further effects of vapor of monobasic ammonium phosphate 188

PLATE 35. Longitudinal section through a normal node and part of the two adjoining internodes of a well-developed plant of *Ricinus communis* 188

PLATE 36. Top of an unopened internode in a stem of *Ricinus communis* next under one that had sealed into it a tube containing 0.2 c. c. of a 20 per cent solution of monobasic ammonium phosphate 188

PLATE 37. 1.—Longitudinal section of a normal young internode of *Ricinus communis*. 2.—Cross section of a Ricinus internode into which was injected a 5 per cent solution of monobasic ammonium phosphate. 3.—Injected internode of another plant in the same series. 4.—Cross section of a normal internode of the same age as that in figure 2 188

PLATE 38. 1.—Same as Plate 37, figure 2, but from an embedded stained section. 2.—A detail of the second xylem-phloem ring showing irregularities more distinctly. 3.—Sieve plate from second xylem-phloem ring. 4.—Distorted sieve tube from third xylem-phloem ring 188

PLATE 39. 1.—Enlarged cross section of a rather regular portion of the second xylem-phloem ring in *Ricinus communis* (Pl. 37, fig. 2). 2.—From the same vascular ring, as figure 1, showing tracheæ lying at right angles to the longer axis of the stem 188

PLATE 40. 1.—Cross section of second xylem-phloem cylinder in Plate 37, figure 2. 2.—Another swirl of cambium on the pith-cavity side of the second xylem-phloem ring in *Ricinus communis* 188

PLATE 41. 1.—Second xylem-phloem ring in Plate 37, figure 2. 2.—A portion of the fourth (imperfect) xylem-phloem ring in Plate 37, figure 2. 3.—Like figure 2, but another part of the fourth xylem-phloem cylinder .. 188

PLATE 42. 1, 2.—Cross sections of an internode of *Ricinus communis*, injected with distilled water containing 5 per cent of dibasic ammonium phosphate 188

PLATE 43. 1.—General view of the small displaced vascular strands shown at *X* in Plate 20. 2.—Structure in cross section of the larger of the two strands in figure 1. 3.—Sieve plates of figure 2 further enlarged 188

PLATE 44. 1.—Cross section of stem of *Ricinus communis* treated with ammonia. 2.—Cross section of Recinus stem treated with strong ammonia 188

PLATE 45. 1.—Inner three-quarters of one of the middle proliferations on Plate 5. 2.—Recinus stem treated with strong ammonia 188

PLATE 46. 1, 2.—Cross sections of the killed tissue (pith) at one side of the proliferations shown on Plate 5. 3.—Cross section of cauliflower leaf, showing intumescences formed after exposure to vapor liberated from one-half gm. of secondary methylamin chloride 188

PLATE 47. 1.—Vertical section of outer part of the proliferations just below the surface shown on Plate 5. 2.—Vertical section in same region as figure 1, but at another level 188

PLATE 48. Portion of under surface of a cauliflower leaf showing intumescences due to vapor of ammonia 188

Page

PLATE 49. 1.—Under surface of a cauliflower leaf from the same series as Plate 48 but nine days after 15 minutes' exposure to vapor of ammonia. 2.—Same series as figure 1, but at the end of 18 days, and from the upper surface of a leaf..... 188

PLATE 50. 1.—Cross section of a normal cauliflower leaf. 2.—Early stage of cauliflower intumescence due to vapor of ammonia..... 188

PLATE 51. 1, 2.—Free-hand unstained sections from old cauliflower intumescences caused by vapor of ammonia..... 188

PLATE 52. 1, 2.—Vertical sections of two ammonia-vapor intumescences on a cauliflower leaf..... 188

PLATE 53. 1.—Like Plate 52 but here the palisade tissue is also involved. 2.—Cross section of an ammonia intumescence above the level of the under surface of the leaf..... 188

PLATE 54. 1, 2.—Cross sections through the deeper tissues of two ammonia intumescences in a cauliflower leaf..... 188

PLATE 55. Cauliflower leaf showing killing effect of mixed ethyl alcohol and acetic acid on maturer leaves of the same plant as Plate 56..... 188

PLATE 56. Under surface of a cauliflower leaf showing intumescences produced by the vapor of mixed ethyl alcohol and acetic acid..... 188

PLATE 57. 1.—Cauliflower intumescences due to vapor of mixed ethyl alcohol and acetic acid. 2.—An enlargement of the right-hand tumor shown in figure 1..... 188

PLATE 58. 1–5.—Cross sections from fixed and stained portions of the little tumors which developed on the leaf shown in Plate 56..... 188

PLATE 59. Cross section of one of the intumescence of Plate 56..... 188

PLATE 60. Vertical sections from fixed and stained portions of cauliflower intumescences shown on Plate 56 and in cross section on Plates 58 and 59. 1.—Middle vascular bundles giving off incipient vessels to the tumor. 2.—Sieve tubes and incipient xylem vessels developing. 3.—Section through the middle of an intumescence which received a *minimum* of stimulation and in which only the epidermis and two layers of cells immediately under it have responded..... 188

PLATE 61. 1.—Like figures 1 and 2 of plate 60, but with the intumescence still covered by an epidermis and free from cork. 2.—Alcohol-acetic-acid intumescence in which entire thickness of the cauliflower leaf is involved..... 188

PLATE 62. 1, 2.—Intumescences formed on a cauliflower leaf exposed 30 minutes in 10½ cubic feet of air space to vapor liberated from one-half gm. of secondary methylamin chlorid. 1.—Under surface of leaf corresponding to figure 2. 2.—Upper surface of same leaf..... 188

PLATE 63. 1.—Same as plate 62, but photographed by transmitted light. 2.—Under surface of one of many leaves of a cauliflower plant exposed to vapor of primary ethylamin..... 188

PLATE 64. 1.—Paraffin-embedded stained section of cauliflower leaf exposed to secondary methylamin. 2. Cross section of cauliflower leaf exposed to primary ethylamin. 3. Same leaf and section as figure 2, but from a normal portion..... 188

PLATE 65. 1.—Portion of ring of concentric medullary bundles in the axis of inflorescence of *Ricinus communis*. 2.—Group of concentric medullary bundles in a branch-gap of *Ricinus communis*..... 188

SOROSPORELLA UVELLA AND ITS OCCURRENCE IN CUTWORMS IN AMERICA

PLATE 66. *Sorosporella uvella:* A.—Larva of *Euxoa tessellata* infected with *Sorosporella uvella*. B, C.—Spore masses. D.—A portion of a spore mass. E.—

Page

a, Spores with ruptured walls of other spores attached. *b*, Spores showing papillæ. F.—Germinating spores. G.—A portion of a spore mass showing peripheral germination. H, K.—Spores germinating, in part showing verticillate branching. L.—Secondary spores. M.—Secondary spores germinating. N.—A portion of vegetative mycelium from a culture. O.—Secondary spores. P.—Secondary spores germinating 194

LEAFSPOT-ROT OF POND LILIES CAUSED BY HELICOSPORIUM NYMPHAEARUM

PLATE 67. A.—Leaf of *Nymphaearum tuberosa* at 1 week after sowing conidia of *Helicosporium nymphaearum* on upper surface. B.—Leaf of *Nymphaea* sp. showing spots due to natural infection with *Helicosporium nymphaearum*. C.—Cross section of young sclerotium of *Helicosporium nymphaearum* from a cornmeal-agar culture. D.—Sclerotium of *H. nymphaearum* from corn-meal-agar culture kept at laboratory temperature for 7 months. E.—Cross section of old sclerotium developed in a corn-meal-agar slant culture 232

PLATE 68. A, B.—Leaves of *Nymphaea caerulea* at 1 week after sowing conidia of *Helicosporium nymphaearum* on upper surface. C.—Leaf of *N. capensis* at 1 week after conidia of *H. nymphaearum* on upper surface 232

PLATE 69. A.—Conidium of *Helicosporium nymphaearum* from culture 3½ months old. B.—Cross section through conidium of *H. nymphaearum*. C.—Conidial group (*H. nymphaearum*) *in situ* in petri-dish culture of corn-meal-agar at 2 weeks. D.—Conidia of *H. nymphaearum*. E.—Conidial group (*H. nymphaearum*) in corn-meal-agar petri-dish culture at 13 days. F.—Water mount of conidia of *H. nymphaearum* from a corn-meal-agar petri-dish culture 13 days old. G.—Conidia of *H. nymphaearum* from corn-meal-agar culture 232

PLATE 70. A.—Cross section of normal part of leaf blade of *Nymphaea* odorata the diseased part of which is shown in B. B.—Cross section of leaf of *N. odorata* at four days after sowing of conidia of *Helicosporium nymphaearum* from pure culture to upper surface of leaf. C.—Cross section of leaf of *N. odorata* at four days after sowing of conidia of *H. nymphaearum* from pure culture to upper surface of leaf 232

STUDIES OF THE GENUS PHYTOPHTHORA

PLATE 71. A.—Mycelium of *P. syringae* grown on potato agar. B.—Mycelium of *P. syringae* grown on oat agar 276

PLATE 72. Conidiophores of the various species of Phytophthora studied from Van Tieghem cells. A.—*P. jatrophae*. B.—*P. arecae*. C.—*P. nicotianae*. D.—*P. phaseoli*. E.—*P. cactorum* (*Panax* sp.). F.—*P. fagi*. G.—*P. cactorum* (*Phyllocactus* sp.). H.—*P. faberi* 276

PLATE 73. Conidia of the various species of Phytophthora illustrating differences in shape of conidia and appearance of the terminal papilla. A.—*P. erythroseptica*. B.—*P. fagi*. C.—*P. arecae*. D.—*P. phaseoli*. E.—*P. parasitica*. F.—*P. infestans* 276

PLATE 74.—Conidia of the various species of Phytophthora illustrating differences in shape of conidia and appearance of the terminal papilla. A.—*P. cactorum* (*Phyllocactus* sp.). B.—*P. syringae*. C.—*P. faberi*. D.—*P. nicotianae*. E.—*P. jatrophae* 276

PLATE 75. Stages in the germination of the conidia and swarm spores of *P. arecae*. A.—Earlier stages. B.—Later stages.

PLATE 76. Chlamydospores and oospores of the various species of Phytophthora, illustrating shape and relation of the antheridium to the oogonium. A.—Oospores of *P. syringae*. B.—Chlamydospores of *P. hatrophae*. C.—

Page

Oospores of *P. arecae*. D.—Chlamydospores of *P. parasitica*. E.—Oospores of *P. erythroseptica*. F.—Oospores of *P. phaseoli*..... 276

PLATE 77. Chlamydospores and oospores of the various species of Phytophthora. A.—Chlamydospores of *P. faberi*. B.—Oospore-like bodies of *P. infestans*. C.—Oospores of *P. cactorum* (*Phyllocactus* sp.). D.—Oospores of *P. fagi*. E.—Chalmydospores of *P. nicotianae*..... 276

TETRASTICHUS BRUCHOPHAGI, A RECENTLY DESCRIBED PARASITE OF BRUCHOPHAGUS FUNEBRIS

PLATE 78. *Tetrastichus bruchophagi:* A.—Larva. B.—Pupa. C.—Adult.... 282

ARSENICAL INJURY THROUGH THE BARK OF FRUIT TREES

PLATE 79. Funnel used in the treatment of trunks with arsenical solutions and suspensions..... 318

PLATE 80. Dissection of bandage used in the treatment of branches with insoluble arsenical compounds..... 318

PLATE 81. Character of rough bark of apple tree: A.—First splits in the formation of rough bark. B.—Roughened condition as found on practically all old trunks and limbs..... 318

PLATE 82. Lenticels of different ages: A.—Lenticels on a young shoot that has just completed the season's growth. B.—Lenticels on a limb about 8 years old. C.—Lenticels on roots..... 318

PLATE 83. A.—Section through a lenticel just forming. B.—Section through a well-developed lenticel. C.—Lenticel spots formed by the absorption of zinc arsenite through lenticels inclosed in arsenical bandages..... 318

PLATE 84. Character of arsenical injury: A.—Branch treated with a solution of arsenic acid. B to I.—Portions of tree injured by a single treatment with sodium-arsenite solution. B.—Cross section of main limb 1 foot below intersection with treated branch. C.—Cross section 16 inches below bandage and just above main limb. D.—Cross section under bandage just below wounds. E.—Cross section 1 foot above bandage. F.—Surface view 8 to 11 inches below bandage. G and H.—Cross sections of side branch that joined treated branch 8 inches below bandage. I.—Main limb with treated branch removed..... 318

A SQUASH DISEASE CAUSED BY CHOANEPHORA CUCURBITARUM

PLATE 85. *Choanephora cucurbitarum:* A.—Immature conidiophore with ramuli developing on the primary vesicle. B.—Budlike processes the beginning of conidial formation on the capitellum. C.—Conidia. D.—Mature capitulum covered with a layer of conidia. E.—Primary vesicle from which ramuli have been detached. F.—Mycelium in tissue of squash-flower pedicel. G.—Germination of conidia 3½ hours old in nutrient agar. H.—Germination of conidia 3½ hours old in water. I.—*a*, Early stage in germination of conidia in hanging drop; *b*, growth of mycelium from same conidia as figure *a* drawn 15 minutes later: *c*, one-half hour later than figure *a*; *d*, 15 minutes later than figure *c*; *e*, one hour later than figure *a*. J.—Germination of sporangiospores. K.—Mycelium of the fungus within the tissues of squash fruit. L.—Variation in size of sporangia and columella. M.—Chlamydospores borne in chains in old cultures. N.—Sporangiospores with tufts of hairlike appendages. O.—Chlamydospores formed singly in 4-day-old cultures..... 328

PLATE 86. *Choanephora cucurbitarum:* A.—Gametes prior to fusion. B.—Young zygospores whose exospore has not yet developed and whose con-

Page

tents are a dense mixture of granular protoplasm and small oil droplets. C.—Lateral view of a young zygospore in the same stage of development as figure B. D.—Cœnocytic mycelium. E.—Mature zygospore with a well-developed exospore and a large oil globule..... 328

PLATE 87. *Choanephora cucurbitarum:* A.—Fructifications on squash corollas and flower pedicels. B.—Natural infection on squash fruit showing the rapidity of the decay when contrasted with figure *C*, which is reproduced from a photograph of the same fruit 15 hours earlier. C.—A diseased teratological squash. D.—Infection through the corolla in the fruit at the left and complete decay of the squash at the right..... 328

DISCOVERY OF INTERNAL TELIA PRODUCED BY A SPECIES OF CRONARTIUM

PLATE 88. A.—Cross section of a petiole of *Ribes roezli*, showing the pericycle (*a*) and pith (*b*) regions which are stuffed with the mycelium of *Cronartium ribicola*. B.—Longitudinal section of the same petiole, showing the manner in which the sori force aside the parenchyma cells of the pith. C.—Internal telium of *Cronartium ribicola* from the pericycle region of a petiole of *Ribes roezli*. D.—Internal telium of *Cronartium ribicola* from the pericycle region of a petiole of *Ribes roezli* flattened by pressure. E.—Young internal telium of *Cronartium ribicola* from the center of the pith region of a petiole of *Ribes roezli*, illustrating the depth at which sori may start to develop..... 332

LITTLE-LEAF OF THE VINE

PLATE 89. A.—A grapevine showing a severe infection of little-leaf. B.—A moderately affected grapevine, showing healthy first leaves..... 398

PLATE 90. A.—Tyloses in the wood of the grapevine. B.—Grapevine shoots showing little leaves and laterals..... 398

PLATE 91 A, B.—Drawings of tyloses in the wood of the grapevine. C, D, E.—Cross section of the trunks of the grapevine. F, G.—Cross sections of healthy and diseased pediocels of the grapevine..... 398

PLATE 92. A.—Grapevines showing the effect of treatment with an injection of copper sulphate. B.—Vineyard showing a severe case of the intensive form of little-leaf..... 398

SPORE-FORMING BACTERIA OF THE APIARY

PLATE 93. A.—*Bacillus mesentericus:* Growth of a 24-hour-old surface colony on an agar plate. B.—*Bacillus vulgatus:* Growth of a 24-hour-old surface colony on an agar plate..... 420

PLATE 94. Growth of 24-hour-old cultures on potato slopes: A.—*Bacillus vulgatus*. B.—*Bacillus mesentericus*. C.—*Bacillus alvei*. D.—*Bacillus orpheus*..... 420

FUSARIUM-BLIGHT, OR WILT DISEASE, OF THE SOYBEAN

PLATE 95. A.—A diseased stem of soybean, showing the roughened appearance caused by the irregular covering of sporodochia. B.—Interior of healthy stem of soybean. C.—Interior of diseased stem of soybean. D.—Soybean plants grown out of doors in the same type of clay soil: *D*, healthy; *E*, diseased through the naturally infected soil..... 440

PEANUT WILT CAUSED BY SCLEROTIUM ROLFSII

PLATE 96. A.—A hill of peanut plants completely wilted by *Sclerotium Rolfsii*. B.—A portion of a wilted shoot showing the white mycelium and sclerotia of *Sclerotium Rolfsii* about its base..... 448

Page

PLATE 97. A.—White mycelium and sclerotia of *Sclerotium Rolfsii* on fallen leaflets. B.—Three hills of peanuts which indicate that *Sclerotium Rolfsii* spreads from wilted plants and attacks adjoining healthy plants..... 448

TEXT FIGURES

DIAGNOSIS OF TUBERCULOSIS BY COMPLEMENT FIXATION, WITH SPECIAL REFERENCE TO BOVINE TUBERCULOSIS

FIG. 1. Specimen of post-mortem report on retained carcasses in Federal meat inspection..... 11

EWES' MILK: ITS FAT CONTENT AND RELATION TO THE GROWTH OF LAMBS

FIG. 1. Curves showing rate of increase in weight of lambs (in pounds) by 4-week periods from different quantities of ewes' milk..... 34

IMMUNITY STUDIES OF ANTHRAX SERUM

FIG. 1. Changes in the serum proteins during the course of hyperimmunization. 51

TEMPERATURE RELATIONS OF APPLE-ROT FUNGI

FIG. 1. Graph showing the development of rot on Ben Davis and York Imperial apples..... 144
2. Graph showing the development of rot on Yellow Newtown apples.. 145
3. Graph showing the development of rot on Winesap apples..... 146
4. Graph showing the development of rot on Winesap apples based on the area instead of the diameter of the rot..... 147
5. Graph showing the development of the rot caused by *Fusarium radicicola* on Yellow Newtown apples..... 148
6. Graph showing the development of the rot caused by *Glomerella cingulata* on Yellow Newtown apples..... 148
7. Graph showing the development of the rot caused by *Neofabraea malicorticis* on Winesap apples..... 148
8. Graph showing the development of the rot caused by *Penicillium expansum* on Winesap apples..... 148
9. Graph showing the development of the rot caused by *Sclerotinia cinerea* (apple) on Winesap apples..... 149
10. Graph showing the development of the rot caused by *Sclerotinia cinerea* (prune) on Winesap apples..... 149
11. Graph showing the development of the rot caused by *Sphaeropsis malorum* on Yellow Newtown apples..... 149
12. Graph showing the development of the rot caused by *Volutella fructi* on Yellow Newtown apples..... 149
13. Graph showing the growth of apple-rot fungi in liquid media..... 154
14. Graph showing the growth of *Alternaria* sp. on corn-meal agar in petri dishes..... 156
15. Graph showing the growth of *Botrytis cinerea* on corn-meal agar in petri dishes..... 156
16. Graph showing the growth of *Cephalothecium roseum* on corn-meal agar in petri dishes..... 156
17. Graph showing the growth of *Fusarium radicicola* on corn-meal agar in petri dishes..... 156
18. Graph showing the growth of *Glomerella cingulata* on corn-meal agar in petri dishes..... 157

Page

19. Graph showing the growth of *Neofabraea malicorticis* on corn-meal agar in petri dishes 157
20. Graph showing the growth of *Penicillium expansum* on corn-meal agar in petri dishes 157
21. Graph showing the growth of *Sclerotinia cinerea* on corn-meal agar in petri dishes 157
22. Graph showing the growth of *Sphaeropsis malorum* on corn-meal agar in petri dishes 158
23. Graph showing the growth of *Volutella fructi* on corn-meal agar in petri dishes 158
24. Graph showing the comparative growth of the different rot fungi on corn-meal agar petri dishes, first experiment 159
25. Graph showing the comparative growth of the different rot fungi on corn-meal agar petri dishes, second experiment 160

SOROSPORELLA UVELLA AND ITS OCCURRENCE IN CUTWORMS IN AMERICA

FIG. 1. *Sorosporella agrotidis* (after Sorokin) 189

MEASUREMENT OF THE INACTIVE, OR UNFREE, MOISTURE IN THE SOIL BY MEANS OF THE DILATOMETER METHOD

FIG. 1. The dilatometer 198

STUDIES OF THE GENUS PHYTOPHTHORA

FIG. 1. Graph showing the variation in length of conidia of *Phytophthora infestans*, *P. fagi*, *P. parasitica*, and *P. jatrophae* 243
2. Graph showing the variation in length of conidia of *Phytophthora phaseoli*, *P. cactorum* (*Phyllocactus* sp.), *P. faberi*, and *P. erythroseptica* 245
3. Graph showing the variation in length of conidia of *Phytophthora nicotianae*, *P. cactorum* (*Panax* sp.) *P. arecae*, and *P. syringae* 249
4. Graph showing the variation in width of conidia of *Phytophthora fagi*, *P. phaseoli*, *P. faberi*, and *P. jatrophae* 250
5. Graph showing the variation in width of *Phytophthora infestans*, *P. cactorum* (*Phyllocactus* sp.), *P. erythroseptica*, and *P. arecae* 251
6. Graph showing the variation in width of *Phytophthora parasitica*, *P. syringae*, *P. cactorum* (*Panax* sp.), and *P. nicotianae* 252
7. Diagrammatic illustration showing that the ratio of the long to the short diameter of ellipsoidal and spherical bodies offers a quantitative means for designating their shape 252
8. Diagram of the arrangement in classes of the ratios of the length to the width of the various conidia, showing the limits of variation and modes 253
9. Camera-lucida drawings showing differences in the shape of the terminal papilla 254
10. Graph showing the variation in the diameters of chlamydospores of *Phytophthora nicotianae*, *P. parasitica*, *P. faberi*, and *P. jatrophae* 257
11. Graph showing the variation in the diameters of oospores of *Phytophthora phaseoli*, *P. arecae*, and *P. erythroseptica* 258
12. Graph showing the variation in the diameters of oospores of *Phytophthora cactorum* (*Phyllocactus* sp.), *P. cactorum* (*Panax* sp.), *P. fagi*, and *P. syringae* 259
13. Graph showing the variation in the diameter measurements of the oospore-like bodies of *Phytophthora infestans* 265

EFFECT OF IRRIGATION WATER AND MANURE ON THE NITRATES AND TOTAL SOLUBLE SALTS OF THE SOIL

Page

FIG. 1. Graph showing the total soluble salts and nitrates in soil held in the laboratory, with different quantities of moisture, for 2½ years 336
2. Graph showing the total soluble salts and nitrates in cropped and fallow soil maintained at various moisture contents in large tanks 337
3. Graph showing the total soluble salts, nitrates, and crop yields in tanks containing soil with varying moisture contents during different stages in the growth of the crop 338
4. Arrangement and treatment of plots in field experiment 339
5. Graph showing the soil moisture and nitrates on August 7, 1911, to a depth of 3 feet in cropped soil to which different quantities of water and manure had been added 340
6. Graph showing the soil moisture and nitrates on August 7, 1911, to a depth of 3 feet in fallow soil to which different quantities of water and manure had been applied 341
7. Graph showing the soil moisture and nitrates on July 21, 1913, to a depth of 10 feet in cropped soil to which different quantities of water and manure had been added 342
8. Graph showing the soil moisture and nitrates on July 21, 1913, to a depth of 10 feet in fallow soil to which different quantities of water and manure had been added 343
9. Graph showing the total soluble salts in the fall in cropped soil receiving different quantities of manure and irrigation water 344
10. Graph showing the total soluble salts in the fall in fallow soil receiving different quantities of manure and irrigation water 346
11. Graph showing the nitrates in the fall in cropped soil receiving different quantities of manure and irrigation water 347
12. Graph showing the nitrates in the fall in fallow soil receiving different quantities of manure and irrigation water 350
13. Graph showing effect of irrigation water on nitrates, total soluble salts, and percentage of moisture in cropped soil at the close of the cropping season 352
14. Graph showing effect of irrigation water on nitrates, total soluble salts, and percentage of moisture in fallow soil in the fall 353
15. Graph showing effect of manure on nitrates, total soluble salts, and percentage of moisture cropped and fallow soil at the close of the cropping season 353
16. Graph showing nitrates, total soluble salts, and moisture at different depths in a cropped soil at the close of the cropping season 354
17. Graph showing nitrates, total soluble salts, and moisture at different depths in a fallow soil in the fall 354
18. Graph showing the total yield of corn stover and grain produced with different irrigation and manuring treatments 357

THE RELATIVE INFLUENCE OF MICROORGANISMS AND PLANT ENZYMS ON THE FERMENTATION OF CORN SILAGE

FIG. 1. Curve showing the disappearance of sugars in series 1 367
2. Curve showing the development of acidity in series 1 368
3. Curve showing the increase in amino nitrogen (and the rate of hydrolysis of protein) in series 1 369
4. Curves showing the formation of alcohol in series 1 and 2 370
5. Curves showing the development of acidity in series 2 371

Page

6. Curve showing the disappearance of sugars in series 2 371
7. Curves showing the increase in amino nitrogen and ammonia nitrogen in series 2 372
8. Curves showing the formation of alcohol in series 3 and 4 373
9. Curves showing the development of acidity in series 4 374
10. Curve showing the increase in amino nitrogen in series 4 375
11. Curves showing the development of acidity in series 5 376
12. Curve showing the formation of alcohol in series 5 377
13. Curves showing the rate of evolution of carbon dioxid. (Curves 2 and 3 are coincident during the first one-half day.) 378

LITTLE-LEAF OF THE VINE

FIG. 1. Diagram of the distribution of grapevines affected with little-leaf: *A*, Moderately affected vineyard; *B*, a very badly affected vineyard ... 383
2. Graph showing the results of application of gypsum for the control of little-leaf 396

SPORE-FORMING BACTERIA OF THE APIARY

FIG. 1. *Bacillus mesentericus:* Smear from an 18-hour-old agar slope 411
2. *Bacillus vulgatus:* Smear from a 9-day-old agar culture 412
3. *Bacillus orpheus:* Smear from a 4-day-old agar culture 413
4. *Bacillus alvei:* Smear from a 48-hour-old agar culture 414
5. *Bacillus larvae:* Smear direct from a diseased larva 416
6. *Bacillus pluton:* Smear direct from a diseased larva 417

FUSARIUM-BLIGHT, OR WILT DISEASE, OF THE SOYBEAN

FIG. 1. *A–G*, Types of macroconidia of the species of Fusarium on soybean. *H*, Cross section of the xylem portion of a diseased soybean stem, showing the invasion of the medullary rays (*a*) and the xylem vessels (*b*) by mycelia of the species of Fusarium on soybean 426

DISSEMINATION OF THE ANGULAR LEAFSPOT OF COTTON

FIG. 1. Diagram of the experimental field of cotton: *A*, Portion of the field in which the row inoculated on May 26, 1916, was located. *B*, Part of the plants diagramed in *A*, showing the reversal of spread between June 27 and July 30. *C*, Top row: Row selected for inoculation on June 30. Second row; same on July 15, showing the slight infection. Below: Chart of neighboring plants showing spread of the disease on July 30 to the west and the relative freedom of the rows to the east .. 467
2. Graphs showing direction of wind and its velocity and the amount of rainfall during thunderstorms 470